KILLS OVER
250
WEED TYPES
AS LISTED

GUARANTEED RESULTS†

Spectracide®

WEED STOP®

for Southern Lawns

HERBICIDA PARA CÉSPEDES DEL SUR

- SEE RESULTS IN 8 HOURS
- WON'T HARM LAWNS*
- KILLS OVER 250 WEED TYPES

AS LISTED, INCLUDING DANDELION, CLOVER & YELLOW NUTSEDGE

ACTIVE INGREDIENTS:	
2,4-D, dimethylamine salt	0.342%
Mecoprop-p, dimethylamine salt	0.123%
Dicamba, dimethylamine salt	0.032%
Sulfentrazone	0.008%
OTHER INGREDIENTS:	99.495%
TOTAL	100.000%

CAS Numbers: 2,4-D dimethylamine salt (2008-39-1), Mecoprop-p dimethylamine salt (66423-09-4), Dicamba dimethylamine salt (2300-66-5), Sulfentrazone (122836-35-5)

†For terms of guarantee, see back panel.

THIS PRODUCT CONTAINS (on weight basis): 0.285% 2,4-D acid equivalent, 0.102% Mecoprop-p acid equivalent, 0.027% Dicamba acid equivalent, Isomer Specific by AOAC Method.

KEEP OUT OF REACH OF CHILDREN

CAUTION See back booklet for additional Precautionary Statements.

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

PRECAUCIÓN Consulte el folleto en la parte posterior para conocer las precauciones adicionales.

Net 1 gal
(128 fl oz / 3.7 L)

READY TO USE™

17-12565

Spectracide®

WEED STOP®

for Southern Lawns

HERBICIDA PARA CÉSPEDES DEL SUR

PRODUCT FACTS

WHAT IT DOES:
Kills all major broadleaf weeds including dandelion, chickweed and clover. Rainproof in as little as 3 hours — won't wash away

WHEN TO USE:
Apply when daytime temperatures are between 50°F and 85°F

WHERE TO USE:
Lawns: Kills the weeds not the lawn (note: Do not use on 'Floratam' or 'Bitterblue' varieties of St. Augustinegrass)

QUESTIONS & COMMENTS:
Call 1-800-917-5438 or visit our website at www.spectracide.com

†GUARANTEED RESULTS OR YOUR MONEY BACK

If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.

*When used as directed. Do not use on 'Floratam' or 'Bitterblue' varieties of St. Augustinegrass.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

NOTA: El comprador asume toda la responsabilidad en caso de no seguir las instrucciones sobre seguridad y modo de empleo.

MADE IN THE USA

Distributed by Spectrum Group
Div. of United Industries Corp.
PO Box 142642, St. Louis, MO 63114-0642

PROOF OF PURCHASE

FPO UPC
0 71121 95711 5

EPA Reg. No. 2217-876-8845 EPA Est. Nos. 39754-WI-1 (R), 9688-MO-1 (U), 79533-WI-1 (BT), 39609-MO-1 (BM) Circled letter is first letter of lot number. 17-12564 © 2007 UIC

ABRA AQUÍ **PEEL HERE**

STOP **READ THE ENTIRE LABEL FIRST.**
ALTO. LEA PRIMERO TODA LA ETIQUETA.

BROADLEAF WEEDS CONTROLLED/MALEZAS DE HOJA ANCHA CONTROLADAS:

Amaranth, purple; Amaranth, slender; Aster, heath; Aster, purple; Aster, white heath; Aster, white prairie; Artichoke; Austrian fieldcress; Bedstraw, catchweed; Bedstraw, smooth; Bedstraw, northern; Bedstraw, pineywoods; Beggarticks, hairy; Betony, Florida; Bindweed, field; Bindweed, hedge; Bitter wintercress; Bittercress, hairy; Bitterweed; Black-eyed Susan; Black medic; Blood flower milkweed; Blue lettuce; Brassbuttons; Bristly oxtongue; Broomweed; Buckhorn; Bullnettle; Burclover; Burdock; Burweed, lawn (spurweed); Buttercup, bulbous; Buttercup, creeping; Buttonweed, common (poorjoe); Buttonweed, Virginia; Carolina geranium (wild); Carpetweed; Catnip; Catsear, spotted (false dandelion); Chamber bitter; Chickweed, common; Chickweed, mouseear; Chickweed, sticky; Chicory; Cinquefoil; Clover, alyce; Clover, crimson; Clover, hop; Clover, rabbit foot; Clover, red; Clover, strawberry; Clover, yellow sweet; Clover, white sweet; Clover, white; Cockle; Cocklebur; Cocklebur, oriental; Cocklebur, spiny; Common mullein; Corn spurry; Creeping beggarweed; Creeping jenny (creeping charlie); Cupid shaving brush; Cudweed; Daisy, English; Daisy, oxeye; Dandelion, common; Dandelion, false; Day flower; Deadnettle; Dichondra; Dock, broadleaf; Dock, curly; Dogbane; Dogfennel; Dollarweed (pennywort); Doveweed; Elderberry; Falseflax; False sunflower; Fiddleneck; Filaree, redstem; Filaree, whitestem; Fleabane, daisy (annual); Fleabane, rough; Galinsoga, hairy; Galinsoga, smallflower; Goldenrod; Ground ivy; Groundsel; Gumweed; Hairy fleabane; Hawkweed, meadow; Hawkweed, mouseear; Hawkweed, orange; Hawkweed, yellow; Healall; Heartleaf drymary; Hemp; Henbit; Hoary cress; Horsenettle; Horseweed; Jimsonweed; Johnny-jumpup violet; Knawel; Knotweed, prostrate; Kochia; Lambsquarters; Lespedeza; Mallow, alkali; Mallow, bristly; Mallow, common; Mallow, dwarf; Mallow, Venice; Marcela; Matchweed; Mexicanweed; Moneywort; Morningglory, bigroot; Morningglory, ivyleaf; Morningglory, Mexican; Morningglory, red; Morningglory, tall; Morningglory, woolly; Mugwort; Mustard, ball; Mustard, black; Mustard, blue; Mustard, hedge; Mustard, Indian; Mustard, tansy; Mustard, tumble; Mustard, white; Mustard, wormseed; Nettle, stinging (burning); Nettle, tall; Nettle, wood; Parsley-piert; Parsnip; Pearlwort; Pennycress (Frenchweed); Pepperweed, perennial; Pepperweed, Virginia (peppergrass); Pigweed, prostate; Pigweed, redroot; Pigweed, Russian; Pigweed, smooth; Pigweed, tumble; Pineappleweed; Plains coreopsis (tickseed); Plantain, blackseed; Plantain, bracted; Plantain, broadleaf; Plantain, buckhorn; Plantain, hoary; Plantain, narrowleaf;

weeds cont. on next page

Plantain, slender; Plantain, woolly; Poison ivy; Poison oak; Pokeweed; Prairie sunflower; Prickly lettuce (compass plant); Prickly sida; Prostrate knotweed; Puncturevine; Purslane, common; Pusley, Brazil; Pusley, Florida; Pusley, large flower; Ragweed, bur; Ragweed, common; Ragweed, lanceleaf; Ragweed, western; Redstem filaree; Rough cinquefoil; Scarlet pimpernel; Shepherdspurse; Smartweed, ladysthumb; Smartweed, pale; Smartweed, Pennsylvania; Smooth chaff-flower; Smooth dock; Sorrel, red (sheep); Sowthistle, annual; Sowthistle, spiny; Spanish needles; Speedwell, bi-lobed; Speedwell, common; Speedwell, corn; Speedwell, ivyleaf; Speedwell, Persian; Speedwell, purslane; Speedwell, slender; Speedwell, snow; Speedwell, thymeleaf; Spiny amaranth; Spurge, spotted; Spurge, prostrate; Spurweed; Star of Bethlehem; Strawberry, Indian mock; Tansy ragwort; Tanweed; Thistle, blessed; Thistle, bull; Thistle, Canada; Thistle, flodman (prairie); Thistle, musk; Thistle, platte; Thistle, plumeless; Thistle, Russian; Thistle, Scotch; Thistle, wavyleaf; Thistle, yellow (horrible); Trailing crownvetch; Velvetleaf; Vervain, blue; Vervain, hoary; Vervain, prostrate; Vervain, tall; Vetch, bird; Vetch, hairy; Vetch, milk; Vetch, narrowleaf; Vetch, wild; Virginia creeper; Western clematis; Western salsify; Wild aster; Wild buckwheat; Wild carrot; Wild four-o-clock; Wild garlic; Wild lettuce; Wild marigold; Wild mustard (yellow rocket); Wild onion; Wild parsnip; Wild radish; Wild rape; Wild strawberry; Wild sweet potato; Wild violet; Woodsorrel, creeping; Woodsorrel, violet; Woodsorrel, yellow (oxalis); Woolly croton; Wormseed; Yarrow, common; Yarrow, western; Yarrow, yellow; Yellow nutsedge

STOP READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Keep people and pets off treated areas until dry.

For home lawn care:
Spectracide® Weed Stop® for Southern Lawns is a ready-to-use product intended for "spot" treatment of individual broadleaf weeds.

- This 1 gal retail container size treats approximately 1,000 weeds or up to 1,000 sq ft.

HOW TO USE

1. Shake well.
2. Connect trigger sprayer. Unscrew and discard the white cap on the bottle. Insert the hose into the bottle and tighten the cap securely.
3. Prime sprayer: Turn the sprayer nozzle one-half turn counterclockwise. Hold the sprayer lower than the bottle and squeeze the trigger several times until it sprays. Adjust the nozzle to coarse spray pattern to minimize wind drift.
4. Spray just enough Spectracide Weed Stop for Southern Lawns to lightly cover broadleaf weeds. ***Excessive spraying can injure some lawns.*** Do not apply when daytime temperatures are above 85°F as lawn injury can occur. Spray only the weeds, not the entire lawn.
5. Weeds will begin wilting and browning overnight.

USE TIPS

- Some hard-to-kill weeds may require re-treatment. Wait at least three weeks between applications — some weeds can take that long to die.
- Rainproof — rain or watering three hours after application will not wash away effectiveness. If rainfall occurs within three hours of application, the product should be reapplied.
- Do not allow spray to drift onto desirable plants since injury may result.
- **Do not use on 'Floratam' or 'Bitterblue' varieties of St. Augustinegrass.**

ALTO LEA PRIMERO TODA LA ETIQUETA. CUMPLA TODAS LAS PRECAUCIONES Y SIGA CUIDADOSAMENTE LAS INSTRUCCIONES.

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta.

Mantenga a las personas y a las mascotas fuera de las áreas tratadas hasta que el producto se seque.

Para cuidar del césped de su hogar: El herbicida para céspedes del sur de Spectracide® es un producto listo para usar para aplicar en tratamientos "localizados" de malezas individuales de hoja ancha.

- Con este envase de 1 galón se pueden tratar aproximadamente 1,000 malezas o hasta 1,000 pies cuadrados.

CÓMO USAR

1. Agite bien.
2. Conecte el gatillo del rociador. Desenrosque y deseche la tapa blanca de la botella. Introduzca la manguera en la botella y ajuste la tapa de modo que quede segura.
3. Cebe el rociador: Gire la boquilla del rociador media vuelta en sentido contrario al de las agujas del reloj. Sostenga el rociador debajo del nivel de la botella y apriete el gatillo varias veces hasta que salga el rociado. Ajuste la boquilla para descargar un rociado grueso para minimizar el arrastre del rociado.
4. Rocíe con el herbicida para céspedes del sur de Spectracide sólo lo suficiente para cubrir ligeramente las malezas de hoja ancha. ***El exceso de rociado puede dañar algunos tipos de césped***. No aplique cuando las temperaturas diurnas sean superiores a 85°F ya que se puede dañar el césped. Rocíe solo las malezas, no todo el césped.
5. Las malezas comenzarán a marchitarse y adquirir color marrón durante la noche.

CONSEJOS SOBRE EL USO

- Puede ser necesario repetir el tratamiento en algunas malezas difíciles de eliminar. Espere al menos tres semanas entre las aplicaciones; algunas malezas pueden demorar ese tiempo en morir.
- A prueba de lluvia – La lluvia o el riego después de las tres horas de la aplicación no reducen la eficacia. Si llueve dentro de las tres horas de la aplicación, se debe volver a aplicar el producto.
- No permita el arrastre del rociado sobre plantas deseables ya que puede causarles daño.
- **No lo use en las variedades 'Floratam' ni 'Bitterblue' del césped de San Agustín.**

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Turn the nozzle to "OFF" position and store with sprayer above the level of liquid in container to prevent leakage. Store in original container in an area inaccessible to children and pets.

DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals:

CAUTION: Harmful if swallowed. Avoid contact with eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

First Aid	
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-917-5438 for emergency medical treatment advice.	

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Questions & Comments? Call 1-800-917-5438 or visit our website at www.spectracide.com

ALMACENAMIENTO Y ELIMINACIÓN

ALMACENAMIENTO DEL PESTICIDA: Gire la boquilla hasta la posición "OFF" (apagado) y guárdelo con el rociador por encima del nivel de líquido en el envase para evitar derrames. Guarde en su envase original en un lugar cerrado que no sea accesible a los niños ni a las mascotas.

ELIMINACIÓN:

Si está vacío: No vuelva a utilizar este envase. Tírelo a la basura, o puede ofrecerlo para reciclarlo, si existe esa opción.

Si está parcialmente vacío: Llame a su servicio de eliminación de desechos más cercano para obtener instrucciones de eliminación. Nunca tire el producto sin usar por ningún desagüe interior ni exterior.

PRECAUCIONES

Riesgos para los seres humanos y los animales domésticos:

PRECAUCIÓN: Nocivo si se ingiere. Evite el contacto con los ojos o la ropa. Cuando utilice este producto, use camisa de manga larga, pantalones largos, calcetines, zapatos y guantes de goma/hule.

Después de utilizar este producto, enjuague los guantes antes de quitárselos, quítese la ropa y lávela por separado antes de volver a usarla y rápidamente lávese bien las manos y la piel que estuvo expuesta con agua y jabón. Quítese la ropa mojada y báñese en regadera tan pronto como sea posible.

Primeros Auxilios	
Si se ingiere:	• Llame inmediatamente a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento. • Si la persona puede tragar, haga que beba poco a poco un vaso de agua. • No induzca el vómito, a menos que así se lo indique el centro de control de envenenamientos o el médico. • No administre nada a una persona que haya perdido el conocimiento.
Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. También puede llamar al 1-800-917-5438 para obtener información acerca del tratamiento médico de urgencia.	

RIESGOS PARA EL MEDIO AMBIENTE: Este producto es tóxico para los invertebrados acuáticos. El arrastre o el escurrimiento pueden afectar de manera adversa a los invertebrados acuáticos y a las plantas que no sean el objetivo. No aplique directamente al agua. No contamine el agua cuando elimine las aguas de lavado del equipo.

NOTA: El comprador asume toda la responsabilidad en caso de no seguir las instrucciones sobre seguridad y modo de empleo.

¿Preguntas y comentarios? Llame al 1-800-917-5438.

Made in the USA. Distributed by Spectrum Group, Div. of United Industries Corp., PO Box 142642, St Louis, MO 63114-0642
EPA Reg. No. 2217-876-8845 EPA Est. Nos. 39754-WI-1(R), 9688-MO-1(U), 79533-WI-1(BT), 39609-MO-1(BM)
Circled letter is first letter of lot number. 17-12564 ©2007 UIC